No.



THE UNITED SHATES OF AMERICA

TO ALL TO VILOM THESE: PRESENTS SHALL COME:

Syngenta Seeds, Inc.

DICCUS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY PROTECTION THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPUSEMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE USE THE BECLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT CORNEL PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'NP2263'

In Creatimonn Thereof, I have hereunto set my hand and caused the seal of the Hant Insisty Protestion Office to be affixed at the City of Washington, D.C. this twenty-ninth day of April, in the year two thousand and eight.

almo

Commissioner
Plant Variety Protection Office
Agricultural Marketing Service

Calmord T. Schafe

ctary of Agriculture

REPRODUCE LOCALLY, Include form number and da	ate on all reprodu	ctions		Form Approved - OMB No. 0581-0055		
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE			The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Peperwork Reduction Act (PRA) of 1995.			
APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE (Instructions and information collection burden statement on reverse)			Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).			
1. NAME OF OWNER		***	TEMPORARY DESIGNATION OR EXPERIMENTAL NAME	3. VARIETY NAME		
Syngenta Seeds, Inc.			EAFERIWENTAL NAME	NP2263		
4. ADDRESS (Street and No., or R.F.D. No., City,	State, and ZIP Cod	de, and Country)	5. TELEPHONE (include area code)	FOR OFFICIAL USE ONLY		
P.O. Box 959			(763) 593-7333	PVPO NUMBER		
Minneapolis, MN 55440			6. FAX (include area code)	200400063		
			(763) 593-7828	FILING DATE		
7. IF THE OWNER NAMED IS NOT A "PERSON", ORGANIZATION (corporation, partnership, asso		8. IF INCORPORATED, GIVE STATE OF INCORPORATION	9. DATE OF INCORPORATION	January 21, 2004		
Corporation		DE	September 24, 1976	-		
10. NAME AND ADDRESS OF OWNER REPRESE	NTATIVE(S) TO S	ERVE IN THIS APPLICATION. <i>(First</i>	person listed will receive all papers)	F FILING AND EXAMINATION FEES:		
•			•	\$ \$ 3652.00		
Ms. Dana S. Rewoldt				R DATE / /21/04/		
Syngenta Seeds Inc.	500			E CERTIFICATION FEE:		
2369 330th Street, P.O. Box Slater IA 50244	500		 	5 s 768.00		
Stater IA 30244		•		D DATE 3/1/08		
				3/1/00		
11. TELEPHONE (Include area code)	12. FAX (Includ		13. E-MAIL			
515-685-5201	515-865	-50/2	dana.rewoldt@syng			
14. CROP KIND (Common Name)	16, FAMILY NA	AME (Botanical)		IN ANY TRANSGENES? (OPTIONAL)		
Corn (dent)	Gramineae		YES NO	COLONIED LIEDA ADUIO DECEDENCE NI IMPERSOR THE		
15. GENUS AND SPECIES NAME OF CROP		RIETY A FIRST GENERATION HYBR	ID? IP SO, PLEASE GIVE THE APPROVED PETITION TO D	SSIGNED USDA-APHIS REFERENCE NUMBER FOR THE DEREGULATE THE GENETICALLY MODIFIED PLANT FOR		
Zea mays L.	. TYES	☑ NO	COMMERICALIZATION			
19. CHECK APPROPRIATE BOX FOR EACH ATTA (Follow instructions on reverse)	ACHMENT SUBMI	TTED	OF CERTIFIED SEED? (See	/ THAT SEED OF THIS VARIETY BE SOLD AS A CLASS Section 83(a) of the Plant Variety Protection Act)		
 Exhibit A. Origin and Breeding History 	of the Variety		21 DOES THE OWNER SPECIF	tems 21 and 22 below) L NO (If "no", go to item 23) / THAT SEED OF THIS VARIETY BE LIMITED AS TO		
b. Exhibit B. Statement of Distinctness			NUMBER OF CLASSES?			
c. Exhibit C. Objective Description of Var	iety		☐ YES ☐ NO	·		
d. Exhibit D. Additional Description of the	Variety (Optional)	•	IF YES, WHICH CLASSES?	☐ FOUNDATION ☐ REGISTERED ☐ CERTIFIED / THAT SEED OF THIS VARIETY BE LIMITED AS TO		
e. v Exhibit E. Statement of the Basis of th	e Owner's Owners	hip	NUMBER OF GENERATIONS			
f. Voucher Sample (2,500 viable untreate verification that tissue culture will be de			YES NO	TD 4.0.0 TOD FACIL CLASS		
repository)	ando naveble te st	resource of the United	IF YES, SPECIFY THE NUMBER 1,2,3, etc. FOR EACH CLASS.			
g. Filing and Examination Fee (\$3,652), n States" (Mail to the Plant Variety Protect		reasurer of the Offited	FOUNDATION REGISTERED CERTIFIED			
20 LIAO TUE MADIETY (MOLUDING ANY HADIVE	CTCD MATERIAL)	OD 4 UVDBID DRODUCED	(If additional explanation is necessary, please use the space indicated on the reverse.) 24. IS THE VARIETY OR ANY COMPONENT OF THE VARIETY PROTECTED BY			
23. HAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) OR A HYBRID PRODUCED FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERRED, OR USED IN THE U. S. OR OTHER COUNTRIES?			INTELLECTUAL PROPERTY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)?			
YES NO			YES NO			
IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TRANSFER, OR USE FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please use space indicated on reverse.)			IF YES, PLEASE GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED REFERENCE NUMBER. (Please use space indicated on reverse.)			
25. The owners declare that a viable sample of basic seed of the variety has been furnished with application and will be replenished upon request in accordance with such regulations as may be applicable, or for a tuber propagated variety a tissue culture will be deposited in a public repository and maintained for the duration of the certificate.						
The undersigned owner(s) is(are) the owner of this sexually reproduced or tuber propagated plant variety, and believe(s) that the variety is new, distinct, uniform, and stable as required in Section 42, and is entitled to protection under the provisions of Section 42 of the Plant Variety Protection Act.						
Owner(s) is (are) informed that false representation herein can jeopardize protection and result in penalties.						
SIGNATORE OF OWNER			SIGNATURE OF OWNER			
NAME (Please print or type)			NAME (Please print or type)			
Ronald S. Ferriss, Ph. D.			Ronald S. Ferriss, Ph. D.			
CAPACITY OR TITLE DATE C.			CAPACITY OR TITLE	DATE		
Director, Strategy Facilitation	01	/09/2004	Director, Strategy Facilitation 01/09/2004			

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 filing fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and provide evidence that name has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, 10301 Baltimore Avenue, Suite 401 NAL Building, Beltsville, MD 20705. Telephone: (301) 504-5682 http://www.ams.usda.gov/lsg/seed.htm.

ITEM

- 19a. Give:
- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

Released in a hybrid, 1-22-2003, offered for sale in the U.S.A. and Canada

24. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NP911 has been issued a PVP Certificate, 1992, #9200012

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filing a change of address; owner's representative; ownership or assignment; or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contect USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.



Syngenta Seeds, Inc. NP2263 Exhibit A

Origin and Breeding History of Corn Inbred Line NP2263

The corn inbred NP2263 was derived from the initial cross of inbred line \$\infty\$911 and inbred line NP903. Inbred Line \$\infty\$911 was developed by self-pollination in Pioneer Brand commercial F1 hybrid 3737. Inbred line NP903 was developed by self-pollination in the Pioneer Brand commercial F1 hybrid 3906.

After the initial cross of 911 and NP903, the breeding method was simple pedigree ear-to-row for the development of inbred line NP2263.

The initial self-pollination of the S0 source was made at Stanton, MN in the Northrup King Company nursery during the summer of 1990. This was followed by self-pollinating and pedigree selection at Stanton, MN and Kekaha, Kauai, Hawaii from 1991 through 1997.

The chronology of inbreeding is given below;

1991 Stanton, MN	Backcross the F ₁ of \$\square\$911 and NP903 to \$\square\$911 to produce S ₀ population.
1991 Kekaha, Kauai	S ₀ population self-pollinated to create S ₁ ears.
1992 Stanton, MN	S ₁ self-pollinated to produce S ₂ ears.
1993 Stanton, MN	Selected S ₂ ears self-pollinated to produce S ₃ ears.
1994 Stanton, MN	Selected S ₃ ear rows self-pollinated to produce S ₄ ears.
1994 Kekaha, Kauai	Selected S ₄ ear rows self-pollinated to produce S ₅ ears.
1995 Stanton, MN	Selected S_5 ear rows self-pollinated to produce S_6 ears.
1995 Kekaha, Kauai	Selected S_6 ear rows self-pollinated to produce S_7 ears.
1996 Stanton, MN	Selected S ₇ ear rows self-pollinated to produce S ₈ ears.
1997 Stanton, MN Seed.	Selected S_8 ear rows self-pollinated and ear rows bulked to produce Breeder

Selection criteria used during the inbreeding process included synchronous male and female flowering, plant health, ear fill, pre-harvest intactness and resistance to various stalk rots. Plants within each generation were also closely evaluated for uniformity of anther and silk color and plant and ear height. Selection was also done for specific combining ability for yield in hybrid combinations across several inbreds.

From 1997 to the present, the inbred line has been observed in London, Ontario, Canada; Stanton, MN; Janesville, WI and other locations. No phenotypic or isozymic variants have been observed. NP2263 is a uniform and stable inbred (from 1997 to 2003 during at least 5 generations of propagation).

Syngenta Seeds, Inc. NP2263 Exhibit B

Distinctness of Corn Inbred Line NP2263

The corn inbred line NP2263 (seed source 2321140) is most similar to the PVP Standard Inbred Line A619 (seed source DY990012). Comparisons of the two varieties were conducted in "side-by-side" trials in 2002 in Stanton, MN (2 plantings) and Janesville, WI (2 plantings) and trials in 2003 in Stanton, MN (2 plantings), Janesville, WI (2 plantings) and London Ont., Canada (1 planting). The trials were planted with two replications at each site. Plot size included 2 rows for each variety and measured 152 cm x 518 cm. Each plot had approximately 70 plants.

For plant level traits, we sampled 5 representative plants from the 2 rows of the 2 row plot at each location/environment and calculated an average for the plot. For plot level traits we evaluated the 2 row plot and gave an average score for the plot within that location/environment.

NP2263 differs from A619 for several different traits. These traits are:

The NP2263 leaf color is darker than the A619 leaf. The NP2263 leaf is a dark green (03 or 5GY 3/4) and the A619 leaf is medium green (02 or 5GY 4/4).

The anther color of NP2263 is purple (17 or 5RP 3/6) and A619 is green-yellow (05 or 2.5GY 8/6).

The cob color of NP2263 is red (14 or 5R 4/10) and A619 is white (19 or 2.5Y 8/2).

NP2263 is a distinct and unique inbred line.

FORM -SD-470-28 (9-1-92)

United States Department of Agriculture, Agricultural Marketing Service Science Division, Plant Variety Protection Office National Agricultural Library Building, Room 500 Beltsville MD 20705

OBJECTIVE DESCRIPTION OF VARIETY CORN (Zea mays L.)

Application val	iety Data		Page 1		Standard	Inbred Data		
4 Application Va		of Brace Roots: 1=Absent 2		4=Dark	2 Standard	Inhand Date		
2.0	0 Average Number of Ears per Stalk 0.4 45 1.0 0.0 45							
0.1	1 Average Number of Tillers 0.2 45			45	0.1		0.2	45
11.8	cm Length of	Top Ear Internode	1.8	45	13.8		8.0	45
54.5	cm Ear Height	t (to base of top ear node)	7.2	45	46.9		2.1	45
	cm Plant Heig	ht (to tassel tip)	Standard Deviation 5.6	Sample Size 45	190.9	Standard Devia	ation 13.5	Sample Size 45
4. PLANT:								
NA	ΝA	From 50% silk to harves	t at 25% moisture			NA		NA
NA	NA From 50% silk to optimum edible quality				NA NA		NA	
004	108.9	From 10% to 90% poller				004		101.4
70					69		1214.3	
72					71		1261.8	
3. MATURIT	uthwest 7=Oth Y (In Region B HEAT UNITS	er est Adaptability; show Hea	t Unit formula in "Co	mments" section	n):	DAYS		HEAT UNITS
		ral 3=Northeast 4=Southe	east 5=Southcentral					
				Standard	d Seed Source DY990012			
2.1=Sweet 2=Dent 3=Flint 4=Flour 5=Pop 6=Ornamental 7=Pipecorn								
WF9 1. TYPE: (d	W64A, A5	54, A654, Pa91 ediate types in Comments se	CI66, Fection)	1105, Ky228		Mo15W, Mo16	W, Mo	24W
C103 Oh43	Mo17, Va	102, Va35, A682 71, H99, Va26	White De	nt:	Р	ipecorn:	-	
В37 В73	B37, B76, N192, A6	, H84 79, B73, NC268	W117, W182B	W153R N	Po	opcorn: SG1533, 4722	2, HP30	1, HP7211
Family B14	CM105, A	.632, B64, B68	Oh7, T2			C13, Iowa512	5, P39,	2132
trial dat		S(Use the most similar(in ba	•	rity)of these to nt (Unrelated):	Sv	veet Corn:		
05=Green-Ye	llow	10=Pink-Orange	15=Red & White	20=White	Capped	25=Variega 26=Other (ited (De Describ	e)
02=Medium (03=Dark Gree 04=Very Dark	Green en	07=Yellow 08=Yellow-Orange 09=Salmon	12=Light Red 13=Cherry Red 14=Red	17=Purpl 18=Color 19=White	e less	22=Tan 23=Brown 24=Bronze		
	CES (Use in cor	njunction with Munsell color 06=Pale Yellow		-	escribe #2			
whole numbe	rs by adding lea	er that describes the varietal ading zeroes if necessary. Ced by a '*' are considered ne	Completeness should	be striven for t	o establis	h an adequate	variety	,
P.O. Box 959, Minneapolis, MN 55440 USA				PVP	O Number 2	005	100063	
Address (Stre	et & No., or R.	F.D. No City, State, Zip Co	de and Country)			OFFICIAL US	E	
Syngenta See Address (Stre	et & No., or R.		232	ety Seed Sourc 1140	NP2	263		ary Designation



Appl	licant Va	ariety Data NP2263	Page 2		Standard	Inbred Data A619	
5. L	EAF:		Standard Deviation	Sample Size		Standard Deviation	Sample Size
	008.3	cm Width of Ear Node Leaf	0.8	45	009.1	0.4	45
	077.9	cm Length of Ear Node Leaf	8.0	45	071.6	7.4	45
	005	Number of leaves above top ear	0.8	45	005	0.2	45
	041	degrees Leaf Angle (measure from 2 nd leaf above ear at ar	20.3 hthesis to stalk above le	45 af)	040	18.0	45
	03	Leaf Color (Munsell code 5GY 3/4)			02	(Munsell code 5G)	(4/4)
	4	Leaf Sheath Pubescence (Rate on scal	e from1+none to 9=like	e peach fuzz)	2		
	5	Marginal Waves (Rate on scale from 1	=none to 9=many)		6		
	4	Longitudinal Creases (Rate on scale fr	om 1=none to 9=many)	5		
6.	TASSEL	•					
	09	Number of Primary Lateral Branches	0.9	45	09	1.7	45
	049	Branch Angle from Central Spike	20.6	45	044	16.9	45
	41.8	Cm Tassel Length	2.5	45	42.4	6.2	45
	7	(from top leaf collar to tassel tip) Pollen Shed (Rate on scale from 0=ma	le sterile to 9=heavy sh	ed)	7	•	
	17	Anther Color (Munsell code 5RP 3/6)			05	(Munsell code 2.5	GY 8/6)
	26	Glume Color (Munsell code 5GY 7/6)	see comments		05	(Munsell code 5G)	7/6)
	2	Bar Glumes (Glume Bands): 1=Absen	t 2=Present	i	2		
7a. E	AR (Un	husked Data):	·				
	26	Silk Color (3 days after emergence) (M see comments	unsell code 2.5GY 8/8)	ļ	05	(Munsell code 2.56	GY 8/8)
	05	Fresh Husk Color(25 days after 50% sil	king) (Munsell code 5G	77/6)	05	(Munsell code 5GY	7/6)
	22	Dry Husk Color (65 days after 50 % sill	king) (Munsell code 2.5)	(8/4)	22	(Munsell code 2.5)	(8/4)
	2	Position of Ear at Dry Husk Stage: 1=0	Jpright 2=Horizontal 3=	Pendent	1		
	6	Husk Tightness (Rate on scale from1=v	very loose to 9=very tigl	nt)	6		
		Husk Extension (at harvest): 1=Short 3=Long (8-10 cm beyond ear tip) 4=Ve	ery long (>10 cm)		2		
7b. E		isked Ear Data): cm Ear Length	Standard Deviation 1.4	Sample Size 45	15.2	Standard Deviation 1.9	Sample Size 45
	36.2	mm Ear Diameter at mid-point	2.6	45	44.0	2.4	45
	82.5	gm Ear Weight	15.5	45	122.3	23.7	45
	14	Number of Kernel Rows	0.5	45	14	0.4	45
	2	Kernel Rows: 1=Indistinct 2=Distinct			2		
	2	Row Alignment: 1=Straight 2=Slightly	Curved 3=Spiral		2		
	13.6	cm Shank Length	2.8	45	10.8	0.7	40
	2	Ear Taper: 1=Slight 2=Average 3=Extr	reme		2		
		Application Variety Data				Standard Inbred D	ata
		Note: Use chart on first page to choose	color codes for color tr	aits			

8. KERNEL (Dried):	Applicant Va	ariety Data NP2263	Page 3		Standard	Inbred Data A619	
0.9, 7 mm Kemel Length				Sample Size	iudiu		
04.5 mm Kernel Thickness				·			45
42.1 % Round Kernels (Shape Grade) 10.2 45 30.3 24.8 45 1 Aleurone Color Pattern: 1=Homozygous 2=Segregating 19 Aleurone Color (Munsell code 2.57 8/6) 07 Hard Endosperm Type: 1=Sweet (su1) 2=Extra Sweet (sx2) 3=Normal Starch 4=High Amylose Starch 15=Valvs Starch 6=High Protein 7=High Lysine 8=Super Sweet (se) 9=High Oil 10=Other 25.8 gm Weight per 100 Kernels (unsized sample) 1.9 45 28.6 2.8 45 9. COB 23.2 mm Cob Diameter at mid-point 2.9 45 26.5 5 sunderd Deviation Sample Size 26.5 5 sunderd Deviation Size 26.5 5 sunderd Deviation Sample Size 26.5 5 sunderd Deviation Sample Size 26.5 5 sunderd Deviation Size Size Size Size Size Size Size Size	07.7	mm Kernel Width	0.3	45	08.8		45
1 Aleurone Color Pattern: 1=Homozygous 2=Segregating 19 Aleurone Color (Munsell code 2.57 8/6) 19 Aleurone Color (Munsell code 2.57 8/6) 07 Hard Endosperm Type: 1=Sweet (su1) 2=Extra Sweet (sh2) 3=Normal Starch 4=High Amylose Starch 5=Wasy Starch 6=High Protein 7=High Lysine 8=Super Sweet (se) 9=High Oil 10=Other 25.8 gm Weight per 100 Kernels (unsized sample) 1.9 45 28.6 2.8 45 9. COB 23.2 mm Cob Diameter at mid-point 2.9 45 14 Cob Color (Munsell code 8.8 4/10) 19 (Munsell code 2.57 8/2) 10. DISEASE RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant): Leave blank if not tested: Leave Race or Strain Options blank if polygenic): A. Leaf Blights, Wilts, and Local Infection Diseases Anthracenes Leaf Blight (Coletorichum graminicola) Common Smut (Victidago maydis) 8 Eyespot (Kabatella zeae) Gos's Wilt (Caracopora zeae-maydis) 6 Helmintosporium Leaf Spot (Egiplaris zeicole) Race 3 6 Northern Leaf Blight (Selectorichum graminicola) Common Smut (Victidago maydis) 8 Eyespot (Kabatella zeae) Gray Leaf Spot (Cerscopora zeae-maydis) 9 Estewart's Wilt (Erwinia stewartii) Other (Specify) 8. Systemic Diseases Corn Lethal Necrosis (MCNV and MDMV) Head Smut (Sphacelotheca reiliana) Mazic Cilhorotic Davi Virus (MDMV) Strain Soripam Downy Mildew of Corn (Peranasolensspora soghi) Other (Specify) 6. Stalk Rots Approfiliae Ear and Kernel Rot (Aspergillus flavus) Dipiodia Stalk Rot (Gilberella zeae) Other (Specify) 7. Stalk Pote (Specify) 6. Stalk Rots Approfiliae Ear and Kernel Rot (Aspergillus flavus) Dipiodia Ear Rot (Sciencoarpella maydis) Gilberella Ear Rot (Gilberella zeae) Other (Specify) 7. Stendard Inbred Data	04.5	mm Kernel Thickness	0.5	45	03.7	0.7	45
19 Aleurone Color (Munsell code 2.5Y 8/6) 07 Hard Endosperm Color (Munsell code 2.5Y 8/10) 3 Endosperm Type: 1=Sweet (su1) 2=Extra Sweet (sh2) 3=Normal Starch 4=High Amylose Starch 6-Waxy Starch 6-High Protein 7-High Lysine 8-Super Sweet (sh) 9-High Oil 10-Other 25.8 gm Weight per 100 Kernels (unsized sample) 1.9 45 28.6 2.8 45 9. COB 23.2 mm Cob Diameter at mid-point 2.9 45 28.6 2.8 45 14 Cob Color (Munsell code 5R 4/10) 10. DISEASE' RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant): leave blank if not tested: leave Race or Strain Options blank if polygenic): A Leaf Bight, Wills, and Local Infection Diseases Anthracrose Leaf Bight (Colletotrichum graminicola) Common Rust (Vercinia sorphin) Common Rust (Vercinia sorphin) Common Smut (Ustiliago maydis) 8 Eyespot (Kabatiella zaee) Goss's Will (Clavibacter michiganense spp. Nebraskense) Gray Leaf Spot (Cercspopra zaee-maydis) 6 Hetminthosporium Leaf Spot (Bigolaris maydis) Race Stehem Rust (Vercinia polysion) Chrer (Specify) 8. Systemic Diseases Corn Lethal Necrosis (MCNV and MDMV) Head Smut (Sphaeclotheca reliana) Maize Chlorotic Poard Virus (MCNV) Maize Chlorotic Rotic Virus (MCNV) Strain Sorphan Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) 6 Strain Stalk Rot (Ciberciapelia maydis) Fusarium Stalk Rot (Gibberella zee) Other (Specify) 6 Fusarium Stalk Rot (Gibberella zee) Other (Specify) 6 Fusarium Stalk Rot (Gibberella zee) Other (Specify) 6 Fusarium Stalk Rot (Gibberella zee) Other (Specify)	42.1	% Round Kernels (Shape Grade)	10.2	45	30.3	24.8	45
07 Hard Endosperm Color (Munsell code 2.57 8/10) 3 Endosperm Type: 1=Sweet (su1) 2=Extra Sweet (sh2) 3=Normal Starch 4=High Amylose Starch 5=Waxy Starch 6=High Protein 7=High Lysine 8=Super Sweet (sh9) 9=High Oil 10=Other 25.8 gm Weight per 100 Kernels (unsized sample) 1.9 45 28.6 2.8 45 9. COB 23.2 mm Cob Diameter at mid-point 2.9 45 26.5 2.6 5.4 45 14 Cob Color (Munsell code 5R 4/10) 10 Diameter at mid-point 2.9 45 26.5 2.6 5.5 45 14 Cob Color (Munsell code 5R 4/10) 10 JISEASE RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant): leave blank if not tested: leave Race or Strain Options blank if polygenic): A Leaf Blight (Value Lave Race or Strain Options blank if polygenic): A Leaf Blight (Value Subject) (Abbutise Steep) Common Rust (Value Subject) (Abbutise Steep) Common Rust (Value Subject) (Abbutise Steep) Goss's Wilt (Clavibacter michiganense spp. Nebraskense) Gray Leaf Spot (Cerscopiora zeae-maydis) 6 Helminthosporium Leaf Spot (Bigolaris maydis) Race Southern Leaf Blight (Exercibium turcicum) Race Mixed Southern Leaf Blight (Exercibium turcicum) Race Mixed Southern Leaf Blight (Exercibium turcicum) Race Mixed Southern Rust (Valucinia polypora) Stewarts Wilt (Grwinia stewartir) Secretary Stewarts Wilt (Grwinia stewartir) Secretary Sec	1	Aleurone Color Pattern: 1=Homozy	gous 2=Segregating		1		
3 Endosperm Type: 1=Sweet (su1) 2=Extra Sweet (sh2) 3=Normal Starch 4=High Armylose Starch 5=Waxy Starch 6=High Protein 7=High Lysine 8=Super Sweet (se) 9=High 0il 10=Other 25.8 gm Weight per 100 Kernels (unsized sample) 1.9 45 28.6 2.8 45 9. COB 23.2 mm Cob Diameter at mid-point 2.9 45 26.5 2.6 45 19 (Munsell code SR 4/10) 10. DISEASER RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant): leave blank if not tested: leave Roce or Strain Options blank if polygenic): A. Leaf Blights, Wilks, and Local Infection Diseases Anthracnose Leaf Blight (Colletotrichum graminicola) Common Bust (Puccinia sorghi) Common Smut (Ustlago maydis) 8 Eyespot (Kabatielia zeae) Goss's Wilk (Calvabacter michiganense spp. Nebraskense) Gray Leaf Spot (Cerscopra zeae-maydis) 6 Helminthosportum Leaf Spot (Biploairis zeicole) Race 3 6 Northern Leaf Blight (Epipolairis zeicole) Race 3 6 Northern Leaf Blight (Epipolairis zeicole) Race 3 Southern Rust (Puccinia stewartiii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reliliana) Maize Chlorotic Nottle Virus (MCNV) Maize Dwarf Mosice Virus (MCNV) Maize Dwarf Mosice Virus (MCNV) Maize Dwarf Mosice Virus (MCNV) Maize Chlorotic Nottle Virus (MCNV) Maize Dwarf Mosice Virus (MCNV) Maize Chlorotic Nottle Virus (MCNV) Maize Dwarf Mosice Virus (MCNV) Maize Chlorotic Nottle Virus (MCNV) Maize Dwarf Mosice Virus (MCNV) Maize Chlorotic Nottle Virus (MCNV) Maize Maize Chlorotic Nottle Virus (MCNV) Maize Maize Chlorotic Nottle Virus (Mostle Maize) Diploida Stalk Rot (Gibberella zeae) Other	19	Aleurone Color (Munsell code 2.5Y 8	/6)		19	(Munsell code 2.5)	(8/6)
### ### ### ### ### ### ### ### ### ##	07	Hard Endosperm Color (Munsell code	e 2.5Y 8/10)		06	(Munsell code 2.5	Y 8/8)
9. COB 23.2 mm Cob Diameter at mid-point 2.9 45 26.5 2.6 2.6 45 14 Cob Color (Munsell code 5R 4/10) 19 (Munsell code 2.5Y 8/2) 10. DISEASE RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant): leave blank if not tested: leave Race or Strain Options blank if polygenic): A. Leaf Blights, Wilts, and Local Infection Diseases Anthracnose Leaf Blight (Collatotrichum graminicola) Common Smut (Vasiliago maydis) 3 Eyespot (Kabadeila zeee) Goss's Wilt (Calvabacter michiganense spp. Nebraskense) Gray Leaf Spot (Cerscopra zeee-maydis) 6 Helminthosporium Leaf Spot (Bigholaris zelcole) Race 3 6 Northern Leaf Blight (Bigholaris zelcole) Race 3 6 Northern Leaf Blight (Bigholaris selcole) Race 3 5 Submart Swilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphaeclotheca reilliana) Maize Chlorotte Dwarf Virus (MCDV) Maize Chlorotte Dwarf Virus (MCDV) Maize Chlorotte Virus (MCMV) Maize Dwarf Mosiac Virus (MCMV) Maize Chlorotte Virus (MCMV) Maize Chlorotte Virus (MCMV) Maize Chlorotte Swilt Virus (MCMV) Maize Dwarf Mosiac Virus (MidWV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diploida Ear Rot (Sitencarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeee) Other (Specify) Application Variety Data	3	4=High Amylose Starch 5=Waxy Sta	arch 6=High Protein 7=		3		
23.2 mm Cob Diameter at mid-point 2.9 45 26.5 2.6 45 14 Cob Color (Munsell code SR 4/10) 19 (Munsell code 2.5Y 8/2) 10. DISEASE RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant); leave blank if not tested: leave Race or Strain Options blank if polygenic): A. Leaf Blights, Wilts, and Local Infection Diseases Anthracnose Leaf Blight (Collectorichum graminicola) Common Rust (Pucchia sorghi) Common Rust (Pucchia sorghi) Common Rust (Pucchia sorghi) Common Rust (Pucchia sorghi) Common Rust (Volabacter michiganense spp. Nebraskense) Gray Leaf Spot (Carscopora zeae-maydis) 6 Helminthosporhum Leaf Spot (Bipolans zecole) Race 3 6 Northern Leaf Blight (Exerohilum turcicum) Race Mixed Southern Leaf Blight (Exerohilum turcicum) Race Mixed Southern Rust (Pucchia polysora) Stewart's Wilt (Crivinia stewarti) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Dwarf Virus (MCDV) Maize Dwarf Mosiac Virus (MDMV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data	25.8	gm Weight per 100 Kernels (unsized	sample) 1.9	45	28.6	2.8	45
10. DISEASE RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant): leave blank if not tested: leave Race or Strain Options blank if polygenic): A. Leaf Blights, Wilts, and Local Infection Diseases Anthracnose Leaf Blight (Colletorichum graminicola) Common Rust (Puccinia sorghi) Common Smut (Usitiago maydis) 8 Eyespot (Kabatielia zeae) Goss's Wilt (Clavibacter michiganense spp. Nebraskense) Gray Leaf Spot (Cerscopora zeae-maydis) 6 Helminthosporium Leaf Spot (Bipolaris zeicole) Race 3 6 Northern Leaf Blight (Exeropinium turcium) Race Mixed Southern Leaf Blight (Bipolaris maydis) Race Southern Rust (Puccinia polysora) Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Dwarf Mosiac Virus (MDMV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data		mm Cob Diameter at mid-point		•	26.5		Sample Size 45
leave blank if not tested: leave Race or Strain Options blank if polygenic): A. Leaf Blights, Wilts, and Local Infection Diseases Anthracnose Leaf Blight (Colletotrichum graminicola) Common Rust (Pucchia sorghi) Common Smut (Vutilago maydis) 8 Eyespot (Kabatielia zeae) Goss's Wilt (Clavibacter michiganene spp. Nebraskense) Gray Leaf Spot (Cerscoproa zeae-maydis) 6 Helminthosporlum Leaf Spot (Bipolaris zeicole) Race 3 6 Northern Leaf Blight (Escenbillum turcicum) Race Mixed Southern Rust (Pucchia polysora) Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chiorotic Owarf Virus (MCDV) Maize Chorotic Covarf Virus (MCDV) Maize Chiorotic Covarf Virus (MCDV) Maize Dwarf Mosiac Virus (MDNV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Gibberella zeae) Other (Specify) Application Varlery Data Standard Inbred Data	14	Cob Color (Munsell code 5R 4/10)			19	(Munsell code 2.5)	′ 8/2)
A. Leaf Blights, Wilts, and Local Infection Diseases Anthracnose Leaf Blight (Collectorichum graminicola) Common Smut (Ustilago maydis) 8 Eyespot (Kabatiella zeae) Goss's Wilt (Clavibacter michiganense spp. Nebraskense) Gray Leaf Spot (Cerscopora zeae-maydis) 6 Helminthosporium Leaf Spot (Bipolaris zeicole) Race 3 6 Northern Leaf Blight (Exserohilum turcicum) Race Mixed Southern Rust (Puccinia polysora) Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwart Virus (MCDV) Maize Chlorotic Mottle Virus (MCDV) Maize Chlorotic Specify) C. Stalk Rot Anthracnose Stalk Rot (Collectorichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspengillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Gibberella zeae) Other (Specify) Application Varlety Data Standard Inbred Data							
Common Rust (Puccinia sorghi) Common Smut (Ustilago maydis) 8 Eyespot (Kabatielia zeae) Goss's Wilt (Clavibacter michiganense spp. Nebraskense) Gray Leaf Spot (Cerscopora zeae-maydis) 6 Helminthosporium Leaf Spot (Bipolaris zeicole) Race 3 6 Northern Leaf Bilght (Eserohilum turcicum) Race Mixed Southern Leaf Bilght (Eserohilum turcicum) Race Mixed Southern Rust (Puccinia polysora) Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Dwarf Virus (MCDV) Maize Dwarf Mosiac Virus (MCDV) Maize Dwarf Mosiac Virus (MCDV) Maize Dwarf Mosiac Virus (MDMV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Gibberelia zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data		ghts, Wilts, and Local Infection Diseas	es	polygenic):			
Goss's Wilt (Clavibacter michiganense spp. Nebraskense) Gray Leaf Spot (Cerscopora zeae-maydis) 6 Helminthosporium Leaf Spot (Bipolaris zeicole) Race 3 6 Northern Leaf Bilght (Exserohilum turcicum) Race Mixed Southern Leaf Bilght (Exserohilum turcicum) Race Mixed Southern Rust (Puccinia polysora) Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Dwarf Mosiae Virus (MCMV)	_	Common Rust (<i>Puccinia sorghi</i>) Common Smut (<i>Ustilago maydis</i>)	ani graniincola)		_		
6 Helminthosporium Leaf Spot (Bipolaris zeicole) Race 3 6 Northern Leaf Blight (Exserohilum turcicum) Race Mixed Southern Rust (Puccinia polysora) Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Dwarf Mosiac Virus (MCMV) Maize Dwarf Mosiac Virus (MCMV) Maize Dwarf Mosiac Virus (MCMV) Maize Chlorotic Dwarf Virus (MCMV) Maize Chlorotic Motitle Virus (MCMV) Maize Dwarf Mosiac Virus (MCMV) Maize Dwarf Mosiac Virus (MDMV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data	8	Goss's Wilt (Clavibacter michiganens			/		
Southern Leaf Blight (Bipolaris maydis) Race Southern Rust (Puccinia polysora) Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Dwarf Mosiac Virus (MCMV) Maize Dwarf Mosiac Virus (MDMV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data		Helminthosporium Leaf Spot (Bipolai	ris zeicole) Race 3				
Stewart's Wilt (Erwinia stewartii) Other (Specify) B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosiac Virus (MDMV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data	ь	Southern Leaf Blight (Bipolaris mayo			б		
B. Systemic Diseases Corn Lethal Necrosis (MCMV and MDMV) Head Smut (Sphacelotheca reiliana) Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosiac Virus (MDMV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data		Stewart's Wilt (Erwinia stewartii)					
Head Smut (Sphacelotheca reiliana) Malze Chlorotic Dwarf Virus (MCDV) Maize Dwarf Mosiac Virus (MDMV) Maize Dwarf Mosiac Virus (MDMV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data	B. Systemi						
Maize Chlorotic Dwarf Virus (MCDV) Maize Chlorotic Mottle Virus (MCMV) Maize Dwarf Mosiac Virus (MDMV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Sibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data			MV)		٠		
Maize Dwarf Mosiac Virus (MDMV) Strain Sorgham Downy Mildew of Corn (Peronosclerospora soghi) Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data		Maize Chlorotic Dwarf Virus (MCDV)					
Other (Specify) C. Stalk Rots Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data		Maize Dwarf Mosiac Virus (MDMV) St				Strain	
Anthracnose Stalk Rot (Colletotrichum graminicola) Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data	C. Stalk Ro	Other (Specify)					
Diplodia Stalk Rot (Stenocarpella maydis) Fusarium Stalk Rot (Fusarium moniliforme) Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data			m graminicola)				
Gibberella Stalk Rot (Gibberella zeae) Other (Specify) Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data		Diplodia Stalk Rot (Stenocarpella ma	ydis)				
Ear and Kernel Rots Aspergillus Ear and Kernel Rot (Aspergillus flavus) Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data		Gibberella Stalk Rot (Gibberella zeae					
Diplodia Ear Rot (Stenocarpella maydis) Fusarium Ear and Kernel Rot (Fusarium moniliforme) Gibberella Ear Rot (Gibberella zeae) Other (Specify) Application Variety Data Standard Inbred Data	Ear and Kern						
Fusarium Ear and Kernel Rot (<i>Fusarium moniliforme</i>) Gibberella Ear Rot (<i>Gibberella zeae</i>) Other (Specify) Application Variety Data Standard Inbred Data							
Other (Specify) Application Variety Data Standard Inbred Data		Fusarium Ear and Kernel Rot (Fusario					
		Other (Specify)	***************************************	***************************************			
			es for color traits		Standard 1	Inbred Data	•

·		200400063				
	/ariety Data NP2263 Page 4	Standard Inbred Data A619				
	T RESISTANCE (Rate from 1 (most susceptible) to 9 (most resistant):					
j 16	eave blank if not tested:					
	Banks Grass Mite (Oligonychus pratensis)	Standard Basiletian Canala Sina				
	Corn Earworm Standard Deviation Sample	Size Standard Deviation Sample Size				
İ	Leaf-Feeding					
	Silk-Feeding:					
	mg larval wt.					
	Ear Damage					
	Corn Leaf Aphid (Rhopalosiphum maidis)					
	Corn Sap Beetle (Carpophilus dimidiatus)					
	European Corn Borer (Osstrinia nubilalis)					
6	1 st Generation (Typically Whorl Leaf Feeding)	5				
	2 nd Generation (Typically Leaf Sheath-Collar feeding)					
	Stalk Tunneling:					
	cm tunneled /plant Fall Armyworm					
	Leaf-Feeding					
	Silk-Feeding:					
	mg larval wt.					
	Maize Weevil (Sitophilus zeamaize)					
	Northern Rootworm (Diabrotica barberi)					
	Southern Rootworm (Diabrotica undecimpunctata)					
	Southwestern Corn Borer (Diatraea grandiosella)					
	Leaf-Feeding					
	Stalk Tunneling:					
	cm tunneled/plant					
	Twospotted Spider Mite (<i>Tetranychus</i>					
	urticae) Western Rootworm (<i>Diabrotica virgifera virgifera</i>)					
12 AGRON	OMIC TRAITS:					
12. AGRON		· ·				
NA	Stay Green (at 65 days after anthesis) (rate on scale from 1=worst	NA NA				
٠.	to 9=excellent.)					
1	% Dropped Ears (at 65 days after anthesis)	1				
_						
0	% Pre-anthesis Brittle snapping	0				
•	O/ Due authoric Book Lodeine					
0	% Pre-anthesis Root Lodging	2				
0	% Post-anthesis Root Lodging (at 65 days after anthesis)	1				
U	70 Post-anthesis Root Loughly (at 05 days after anthesis)					
1962.1	Kg/ha Yield of Inbred Per Se (at 12-13% grain moisture)	2061.0				
1002.1	rigi, na riola di Illarda i di Galla Illarda Galla Illarda Galla Illarda Galla Illarda Galla Gal					
13. MOLECU	JLAR MARKERS: (0=data unavailable: 1 data available but not supplied:	2=data supplied)				
	,	,				
1	Isozymes 2 RFLP's RAPD's					
NP2263						
REFERENCE	S:					
	R. 1954. A System for the Classification of corn Inbred Lines. PhD thesis					
	R. A., G.W. Beadle, and A.C. Fraser. 1935. A Summary of Linkage Studie G.F. Bills, G.P. Chamuris, A.Y. Rossman. 1989. Fungi on Plant and Plant					
	opathological Society, St. Paul, MN.	Froducts in the officed States. The American				
Inglett, G.E. (Ed) 1970. Corn: Culture, Processing, Products. Avi Publishing Company, Westport, CT.						
Jugenheimer, R.W. 1976. Corn: Improvement. Seed Production, and Uses. John Wiley & Sons, New York.						
McGee, D.C. 1988. Maize Diseases. AOS Press, St. Paul, MN. 150 pp.						
Munsell Color Chart for Plant Tissues. Mcbeth. P.O. Box 230. Newburgh, N.Y. 12551-0230						
The Mutants of Maize. 1968. Crop Science Society of America. Madison, WI.						
Shurtleff, M.C. 1980. Compendium of Corn Diseases. APS Press, St. Paul, MN. 105 pp.						
Sprague, G.F., and J.W. Dudley (Editors). 1988 Corn and Corn Improvement, Third Edition. Agronomy Monograph 18. ASA, SCCA,						
SSSA,						
Madison, W	Madison, W1. Stringfield, G.H, Maize Inbred Lines of Ohio. Ohio A.E.S., Bul. 831. 1959					
U.S. Department of Agriculture. 1936, 1937. Yearbook.						
	amone or righteditare. 1990, 1997. Tearbook.					

NP2263

COMMENTS (eg. state how heat units were calculated, standard inbred seed source, and /or where data was collected. Continue in Exhibit D):

Heat Units per day were calculated using the standard formula: HU={MaxTemp (86) + Min Temp (50)]/2-50.

Data for this exhibit was collected at London, Ontario, Canada, Stanton, MN and Janesville, WI.

3) Large standard deviations are probably due to environmental factors at each individual location where the variety was observed. Since the varieties reported in this exhibit are inbreds, the response to the environment is probably more pronounced than a hybrid or a combination of these inbred lines. Any stress at specific times during the growing season could influence results.

* - There is a slight pale purple shading of the silk ends on A619 with age.

5) Disease and Insect information for both inbred lines was collected at Stanton, MN in 2002.


6) Glume color of NP2263 is green-yellow (05) with dark purple shading (17).7) Silk color of NP2263 is green-yellow (05) with pale purple shading (16).

The environments had different planting dates and were in different fields; these contribute to the variability of the traits.

The London trial location may be causing the standard deviations for these plants to be a bit wider than normal for a couple of reasons. The planting dates for the London trials were substantially later than the dates for the Stanton planting. London is in an area that plants 83-90 day corn and Stanton and Janesville are locations that plant 94-102 day corn. In addition to the relative maturity difference between the locations, the delayed plantings due to the weather conditions in London provide even less time for the plants to mature.


London normally plants in the May 5- May 10 range. But in both years the actual planting dates were substantially later than this date. The environment in 2003 was hotter than normal. To the extent that London trial resorted to nursery irrigation to attempt to avoid moisture stress at flowering time.

Additional Description of the Corn Inbred Line NP2263 Table 1




Typical A619 Silk and Second Leaf Color

Additional Description of the Corn Inbred Line NP2263 Table 1 (continued)




Typical NP2263 Silk and Second Leaf Color

Additional Description of the Corn Inbred Line NP2263 Table 2




Typical A619 Tassel

Additional Description of the Corn Inbred Line NP2263 Table 2 (continued)




Typical NP2263 Tassel

Additional Description of the Corn Inbred Line NP2263 Table 3



Anther and Glume Color of A619



Anther and Glume Color of NP2263

REPRODUCE LOCALLY. Include form number and edition date on al	Il reproductions.	FORM APPROVED - OMB No. 0581-005			
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). The information is held confidential until the certificate is issued (7 U.S.C. 2426).				
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3, VARIETY NAME			
Syngenta Seeds, Inc.	OR EXPERIMENTAL NUMBER	NP2263			
4.40000000					
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (Include area code)			
P.O. Box 959	(763) 593-7333	(763) 593-7828			
Minneapolis, MN 55440	7. PVPO NUMBER 20	0400063			
8. Does the applicant own all rights to the variety? Mark an "X" in the	e appropriate block. If no, please expla	in. YES NO			
9. Is the applicant (individual or company) a U.S. national or a U.S. b	pased company? If no, give name of c	ountry. YES NO			
10. Is the applicant the original owner?	NO If no, please answer <u>one</u>	of the following:			
a. If the original rights to variety were owned by individual(s), is	(are) the original owner(s) a U.S. Nation NO If no, give name of count				
b. If the original rights to variety were owned by a company(ies) YES	, is (are) the original owner(s) a U.S. ba NO If no, give name of count				
11. Additional explanation on ownership (Trace ownership from original	nal breeder to current owner. Use the re	everse for extra space if needed):			
(See attached Exhibit E Addendum)					
(See attached Extitot I Addendum)					
·					
	•				
PLEASE NOTE:					
Plant variety protection can only be afforded to the owners (not licens	sees) who meet the following criteria:				
If the rights to the variety are owned by the original breeder, that protection accountry which affords similar protection to nationals of a country which affords similar protection to nationals or the similar protection.	erson must be a U.S. national, national of the U.S. for the same genus and speci	of a UPOV member country, or es.			
If the rights to the variety are owned by the company which employ nationals of a UPOV member country, or owned by nationals of a genus and species.	yed the original breeder(s), the company country which affords similar protection t	must be U.S. based, owned by o nationals of the U.S. for the same			
3. If the applicant is an owner who is not the original owner, both the	original owner and the applicant must m	eet one of the above criteria.			
The original breeder/owner may be the individual or company who did Act for definitions.	rected the final breeding. See Section 4	1(a)(2) of the Plant Variety Protection			
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, control number. The valid OMB control number for this information collection is 0581-0055. including the time for reviewing the instructions, searching existing data sources, gathering a	The time required to complete this information collec-	tion is estimated to average 0.1 hour per response.			

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.

Syngenta Seeds, Inc. NP2263 Exhibit E, Block #11

Additional Explanation on Ownership

Syngenta breeder, Christopher Perry, developed inbred NP2263, working primarily from Stanton, MN (see Exhibit A for detail). Perry was a breeder for Northrup King Company at the beginning of the line's development, and was employed by Novartis Seeds, Inc. (formed by the merger of Ciba and Sandoz) at the completion of the lines development in 1999. This line is now exclusively owned by Syngenta Seeds, Inc. (formed by the merger of the agribusinesses of Novartis and Zeneca in 1999).